


# Roots of a Nonlinear Equation

Topic: Bisection Method

Major: Mechanical Engineering


# Basis of Bisection Method

**Theorem:** An equation  $f(x)=0$ , where  $f(x)$  is a real continuous function, has at least one root between  $x_l$  and  $x_u$  if  $f(x_l) f(x_u) < 0$ .





# Theorem

If function  $f(x)$  in  $f(x)=0$  does not change sign between two points, roots may still exist between the two points.




# Theorem


If the function  $f(x)$  in  $f(x)=0$  does not change sign between two points, there may not be any roots between the two points.



# Theorem

If the function  $f(x)$  in  $f(x)=0$  changes sign between two points, more than one root may exist between the two points.






# Algorithm for Bisection Method

# Step 1


- Choose  $x_l$  and  $x_u$  as two guesses for the root such that  $f(x_l) f(x_u) < 0$ , or in other words,  $f(x)$  changes sign between  $x_l$  and  $x_u$ .



## Step 2


Estimate the root,  $x_m$  of the equation  $f(x) = 0$  as the mid-point between  $x_\ell$  and  $x_u$  as

$$x_m = \frac{x_\ell + x_u}{2}$$



# Step 3

Now check the following



If  $f(x_l) f(x_m) < 0$ , then the root lies between  $x_R$  and  $x_m$ ; then  $x_l = x_l$  ;  
 $x_u = x_m$ .

If  $f(x_R) f(x_m) > 0$ , then the root lies between  $x_m$  and  $x_u$ ; then  $x_l = x_m$  ;  
 $x_u = x_u$ .

If  $f(x_l) f(x_m) = 0$ ; then the root is  $x_m$ .  
Stop the algorithm if this is true.



# Step 4

---

New estimate

$$x_m = \frac{x_\ell + x_u}{2}$$


Absolute Relative Approximate Error

$$|\epsilon_a| = \left| \frac{x_m^{new} - x_m^{old}}{x_m^{new}} \right| \times 100$$

$x_m^{old}$  = previous estimate of root

$x_m^{new}$  = current estimate of root

# Step 5






# Example

---

- A trunnion has to be cooled before it is shrink fit into a steel hub




# Solution

The equation that gives the temperature 'x' to which it has to be cooled to obtain the desired contraction is given by:


$$f(x) = -0.50598 \times 10^{-10} x^3 + 0.38292 \times 10^{-7} x^2 + 0.74363 \times 10^{-4} x + 0.88318 \times 10^{-2} = 0$$

Use the bisection method of finding roots of equations to find the temperature to which the trunnion should be cooled down to.




# Graph of function $f(x)$

$$f(x) = -0.50598 \times 10^{-10} x^3 - 0.38292 \times 10^{-7} x^2 + 0.74363 \times 10^{-4} x + 0.88318 \times 10^{-2} = 0$$



# Checking if the bracket is valid



Choose the bracket


$$x_l = -100$$

$$x_u = -150$$

$$f(-100) = 1.82902 \times 10^{-3}$$

$$f(-150) = -1.2903 \times 10^{-3}$$

# Iteration #1



$$x_l = -100, x_u = -150$$

$$x_m = \frac{-100 + (-150)}{2} = -125$$

$$f(-100) = 1.82902 \times 10^{-3}$$


$$f(-150) = -1.2903 \times 10^{-3}$$

$$f(-125) = 2.33562 \times 10^{-4}$$

$$x_l = -125$$

$$x_u = -150$$

# Iteration #2



$$x_l = -125, x_u = -150$$

$$x_m = \frac{-125 + (-150)}{2} = -137.5$$

$$|\epsilon_a| = 9.0909\%$$



$$f(-125) = 2.33562 \times 10^{-4}$$

$$f(-150) = -1.2903 \times 10^{-3}$$

$$f(-137.5) = -5.3762 \times 10^{-4}$$

$$x_l = -125, x_u = -137.5$$

# Iteration #3



$$x_\ell = -125, x_u = -137.5$$

$$x_m = \frac{-125 + (-137.5)}{2} = -131.25$$

$$|\epsilon_a| = 4.7169\%$$

$$f(-125) = 2.33562 \times 10^{-4}$$

$$f(-137.5) = -5.3762 \times 10^{-4}$$

$$f(-131.25) = -1.54303 \times 10^{-4}$$



# Convergence

Table 1: Root of  $f(x)=0$  as function of number of iterations for bisection method.

Iteration	$X_l$	$X_u$	$X_m$	$ \epsilon_a  \%$	$f(x_m)$
1	-100	-150	-125	-----	$2.3356 \times 10^{-4}$
2	-125	-150	-137.5	9.0909	$-5.3762 \times 10^{-4}$
3	-125	-137.5	-131.25	4.7619	$-1.5430 \times 10^{-4}$
4	-125	-131.25	-128.125	2.4390	$3.9065 \times 10^{-5}$
5	-128.125	-131.25	-129.6875	1.2048	$-5.7760 \times 10^{-5}$
6	-128.125	-129.6875	-128.9062	0.6061	$-9.3826 \times 10^{-6}$
7	-128.125	-128.9062	-128.5156	0.3039	$1.4838 \times 10^{-5}$
8	-128.5156	-128.9062	-128.7109	0.1517	$2.7228 \times 10^{-6}$
9	-128.7109	-128.9062	-128.8086	0.0758	$-3.3305 \times 10^{-6}$
10	-128.7109	-128.8086	-128.7597	0.0379	$-3.0396 \times 10^{-7}$



# Advantages

---

- Always convergent
- The root bracket gets halved with each iteration - guaranteed.



# Drawbacks

---

- Slow convergence




## Drawbacks (continued)

---

- If one of the initial guesses is close to the root, the convergence is slower

# Drawbacks (continued)

- If a function  $f(x)$  is such that it just touches the  $x$ -axis it will be unable to find the lower and upper guesses.



# Drawbacks (continued)

- Function changes sign but root does not exist

